Private Duty Nursing Acuity Grid Updated October 2007

PRIVATE DUTY NURSING ACUITY GRID

(To be completed by the person completing the patient care)

Recipient	MID#		· · · · · · · · · · · · · · · · · · ·
ASSESSMENT NEEDS		POINTS	SCORE
(choose one)			
Minimal ongoing assessments (less often than Q 6 hrs; at lea	st daily) □	2.00	
Moderate ongoing assessments (hands-on Q 4-6 hrs) □		3.00	
(choose one if at least 2 of the 4 assessments are ordered	d and documented)		
VS/GLU/NEURO/Resp (Assess less often than Q 4, at least of	daily) □	1.00	
VS/GLU/NEURO/Resp (Assess Q 4 hr or more often) □		2.00	
VS/GLU/NEURO/Resp (Assess Q 2 hr or more often) □		3.00	
	TOTAL		
MEDICATION/IV DELIVERY NEEDS		<u>POINTS</u>	SCORE
(choose one if applicable - does not include nebulizer me	ds)		
Oral or G Tube, NG, NJ:			
Medication delivery 1 to 3 doses per day \Box		1.00	
Medication delivery 4 to 6 doses per day \Box		2.00	
Medication delivery 7 doses per day or more □		4.00	
(choose one)			
No IV access □		0.00	
Peripheral IV access □		1.00	
Central Line of port, PICC Line, Hickman □		2.50	
(choose one)			
No IV medication delivery □		0.00	
Transfusion or IV Tx less than daily but at least weekly □		2.50	
V Tx less often than Q 4 hrs (does not include hep flush) □		4.50	
V Tx Q 4 or more often □		6.00	
(choose any that apply)			
Reg blood draws/IV Peripheral Site (#)		4.5*	
Reg blood draws/IV Central Line (#)		6.0*	
TPN		6.00	
· · · · -	TOTAL	0.00	
	. •		

Home Health Manual page 1 of 4

Utah Medicaid Provider Manual	Private Duty Nursing Acuity Grid
Division of Health Care Financing	Updated October 2007

FEEDING NEEDS	POINTS	SCORE
(choose any that apply)		
Routine oral feeding □	0.00	
Difficult, prolonged oral feeding □	2.00	
Occasional reflux and/or aspiration precautions	0.50	
G-Tube, J-Tube, or Mic-key button □	0.50	
(choose one)		
No tube feeding □	0.00	
Tube feeding (routine bolus or continuous) □	2.00	
Tube feeding (combination bolus and continuous) \Box	2.50	
Complicated tube feeding, residual checks, aspiration precautions,	3.00	
(slow feed or other problems) □		
тот	ΓAL	

TOTAL		
RESPIRATORY NEEDS	POINTS	SCORE
(choose one)		
No trach, patent airway □	0.00	
No trach, unstable airway (desats common, airway clearance issues) □	1.00	
Trach (routine care) □	1.00	
Trach (special care - wounds, breakdown, frequent pull-out, replacement) □	2.50	
(choose one)		
No suctioning □	0.00	
Infrequent suctioning (less than Q 8 but at least daily) □	0.50	
Suctioning Q 3 to Q 8 hrs (#)	1.50	
Suctioning Q 2 hrs or more frequently (#)	2.50	
(choose one)		
Oxygen - daily use □	1.00	
Oxygen PRN based on pulse oximetry, oxygen needed at least weekly	0.50	
Humidification (direct) □	0.50	
(choose one)		
No ventilator □	0.00	
Ventilator; rehab transition/active weaning □	9.00	
Ventilator; weaning achieved □	6.00	
Ventilator; non-invasively at night □	8.00	
Ventilator; less than 12 hrs per day □	10.00	
Ventilator; ≥ 12 hrs per day but not continuous □	12.00	
Ventilator; no respiratory effort or 24 hr/day in assist mode □	14.00	
(choose one)		
No BiPAP or CPAP □	0.00	
BiPAP or CPAP up to 8 hrs per day □	4.00	

page 2 of 4 Home Health Manual

Utah Medicaid Provider Manual	Private Duty Nursing Acuity Grid
Division of Health Care Financing	Updated October 2007

BiPAP or CPAP greater than 8 hrs per day □	6.00	
BiPAP ST (with rate) used to ventilate at night □	7.00	
BiPAP ST (with rate) with trach □	8.00	
(choose one)		
No Nebulizer treatments □	0.00	
Nebulizer treatments less than daily but at least QW: # □	1.00	
Nebulizer treatment Q 4 or less frequently: # □	1.50	
Nebulizer treatment Q 3 hrs: #	2.00	
Nebulizer treatment Q 2 hrs or more frequently: # □	3.0*	
(choose one)		
No Chest PT, ABI vest □	0.00	
Chest PT, ABI vest or Cough Assist/less than daily, at least QW: # □	0.50	
Chest PT, ABI vest or Cough Assist/Q 4 or less frequently: # □	1.50	
Chest PT, ABI vest or Cough Assist/Q 3 hrs: # □	2.00	
Chest PT, ABI vest or Cough Assist/Q 2 hrs or more: # □	3.0*	
TOTAL		
ELIMINATION NEEDS	POINTS	SCORE
(choose those that best describe)		
Uncontrolled incontinence < 3 yrs of age □	0.00	
Continence of bowel and bladder □	0.00	
Uncontrolled incontinence, either bowel or bladder, > 3 yrs of age □	1.00	
Uncontrolled incontinence, both bowel and bladder, ≥ 3 yrs of age □	2.00	
_ :		
Intermittent straight catheter □	3.50	
Intermittent straight catheter \Box Uncontrolled incontinence (frequent linen change), \geq 3 yrs of age \Box	3.50 6.00	
Intermittent straight catheter \square Uncontrolled incontinence (frequent linen change), ≥ 3 yrs of age \square Ostomy care - at least daily \square	3.50	
Intermittent straight catheter □ Uncontrolled incontinence (frequent linen change), ≥ 3 yrs of age □ Ostomy care - at least daily □ TOTAL	3.50 6.00 3.00	
Intermittent straight catheter □ Uncontrolled incontinence (frequent linen change), ≥ 3 yrs of age □ Ostomy care - at least daily □ TOTAL SEIZURES	3.50 6.00	SCORE
Intermittent straight catheter □ Uncontrolled incontinence (frequent linen change), ≥ 3 yrs of age □ Ostomy care - at least daily □ TOTAL SEIZURES (choose one)	3.50 6.00 3.00 POINTS	SCORE
Intermittent straight catheter □ Uncontrolled incontinence (frequent linen change), ≥ 3 yrs of age □ Ostomy care - at least daily □ TOTAL SEIZURES (choose one) No seizure activity □	3.50 6.00 3.00 POINTS	SCORE
Intermittent straight catheter □ Uncontrolled incontinence (frequent linen change), ≥ 3 yrs of age □ Ostomy care - at least daily □ TOTAL SEIZURES (choose one)	3.50 6.00 3.00 POINTS 0.00 0.00	SCORE
Intermittent straight catheter □ Uncontrolled incontinence (frequent linen change), ≥ 3 yrs of age □ Ostomy care - at least daily □ TOTAL SEIZURES (choose one) No seizure activity □	3.50 6.00 3.00 POINTS	SCORE
Intermittent straight catheter □ Uncontrolled incontinence (frequent linen change), ≥ 3 yrs of age □ Ostomy care - at least daily □ TOTAL SEIZURES (choose one) No seizure activity □ Mild seizures - at least daily, no intervention □	3.50 6.00 3.00 POINTS 0.00 0.00	SCORE
Intermittent straight catheter □ Uncontrolled incontinence (frequent linen change), ≥ 3 yrs of age □ Ostomy care - at least daily □ TOTAL SEIZURES (choose one) No seizure activity □ Mild seizures - at least daily, no intervention □ Mod seizures (req min intervention - at least daily) □	3.50 6.00 3.00 POINTS 0.00 0.00 2.00	SCORE
Intermittent straight catheter □ Uncontrolled incontinence (frequent linen change), ≥ 3 yrs of age □ Ostomy care - at least daily □ TOTAL SEIZURES (choose one) No seizure activity □ Mild seizures - at least daily, no intervention □ Mod seizures (req min intervention - at least daily) □ Mod seizures (req min intervention - 2 to 4 times per day) □	3.50 6.00 3.00 POINTS 0.00 0.00 2.00 4.00	SCORE
Intermittent straight catheter □ Uncontrolled incontinence (frequent linen change), ≥ 3 yrs of age □ Ostomy care - at least daily □ TOTAL SEIZURES (choose one) No seizure activity □ Mild seizures - at least daily, no intervention □ Mod seizures (req min intervention - at least daily) □ Mod seizures (req min intervention - 2 to 4 times per day) □ Mod seizures (req min intervention - ≥ 5 times per day) □	3.50 6.00 3.00 POINTS 0.00 0.00 2.00 4.00 4.50	SCORE
Intermittent straight catheter □ Uncontrolled incontinence (frequent linen change), ≥ 3 yrs of age □ Ostomy care - at least daily □ TOTAL SEIZURES (choose one) No seizure activity □ Mild seizures - at least daily, no intervention □ Mod seizures (req min intervention - at least daily) □ Mod seizures (req min intervention - 2 to 4 times per day) □ Mod seizures (req min intervention - ≥ 5 times per day) □ Severe seizures (req IM/IV/Rectal med administration - at least daily) □ Severe seizures (req IM/IV/Rectal med administration - 2 to 4 times per day) □	3.50 6.00 3.00 POINTS 0.00 0.00 2.00 4.00 4.50 5.00	SCORE
Intermittent straight catheter □ Uncontrolled incontinence (frequent linen change), ≥ 3 yrs of age □ Ostomy care - at least daily □ TOTAL SEIZURES (choose one) No seizure activity □ Mild seizures - at least daily, no intervention □ Mod seizures (req min intervention - at least daily) □ Mod seizures (req min intervention - 2 to 4 times per day) □ Mod seizures (req min intervention - ≥ 5 times per day) □ Severe seizures (req IM/IV/Rectal med administration - at least daily) □	3.50 6.00 3.00 POINTS 0.00 0.00 2.00 4.00 4.50 5.00 5.50	SCORE
Intermittent straight catheter □ Uncontrolled incontinence (frequent linen change), ≥ 3 yrs of age □ Ostomy care - at least daily □ TOTAL SEIZURES (choose one) No seizure activity □ Mild seizures - at least daily, no intervention □ Mod seizures (req min intervention - at least daily) □ Mod seizures (req min intervention - 2 to 4 times per day) □ Mod seizures (req min intervention - ≥ 5 times per day) □ Severe seizures (req IM/IV/Rectal med administration - at least daily) □ Severe seizures (req IM/IV/Rectal med administration - 2 to 4 times per day) □ Severe seizures (req IM/IV/Rectal med administration - 2 to 4 times per day) □ TOTAL	3.50 6.00 3.00 POINTS 0.00 0.00 2.00 4.00 4.50 5.00 5.50 6.00	
Intermittent straight catheter □ Uncontrolled incontinence (frequent linen change), ≥ 3 yrs of age □ Ostomy care - at least daily □ TOTAL SEIZURES (choose one) No seizure activity □ Mild seizures - at least daily, no intervention □ Mod seizures (req min intervention - at least daily) □ Mod seizures (req min intervention - 2 to 4 times per day) □ Mod seizures (req min intervention - ≥ 5 times per day) □ Severe seizures (req IM/IV/Rectal med administration - at least daily) □ Severe seizures (req IM/IV/Rectal med administration - 2 to 4 times per day) □ Severe seizures (req IM/IV/Rectal med administration - ≥ 5 times per day) □ TOTAL THERAPIES/ORTHOTICS/CASTING	3.50 6.00 3.00 POINTS 0.00 0.00 2.00 4.00 4.50 5.00 5.50	SCORE
Intermittent straight catheter □ Uncontrolled incontinence (frequent linen change), ≥ 3 yrs of age □ Ostomy care - at least daily □ TOTAL SEIZURES (choose one) No seizure activity □ Mild seizures - at least daily, no intervention □ Mod seizures (req min intervention - at least daily) □ Mod seizures (req min intervention - 2 to 4 times per day) □ Mod seizures (req min intervention - ≥ 5 times per day) □ Severe seizures (req IM/IV/Rectal med administration - at least daily) □ Severe seizures (req IM/IV/Rectal med administration - 2 to 4 times per day) □ Severe seizures (req IM/IV/Rectal med administration - ≥ 5 times per day) □ TOTAL THERAPIES/ORTHOTICS/CASTING (choose any that apply)	3.50 6.00 3.00 POINTS 0.00 0.00 2.00 4.00 4.50 5.00 5.50 6.00	
Intermittent straight catheter □ Uncontrolled incontinence (frequent linen change), ≥ 3 yrs of age □ Ostomy care - at least daily □ TOTAL SEIZURES (choose one) No seizure activity □ Mild seizures - at least daily, no intervention □ Mod seizures (req min intervention - at least daily) □ Mod seizures (req min intervention - 2 to 4 times per day) □ Mod seizures (req min intervention - ≥ 5 times per day) □ Severe seizures (req IM/IV/Rectal med administration - at least daily) □ Severe seizures (req IM/IV/Rectal med administration - 2 to 4 times per day) □ Severe seizures (req IM/IV/Rectal med administration - ≥ 5 times per day) □ TOTAL THERAPIES/ORTHOTICS/CASTING (choose any that apply) Fractured or casted limb □	3.50 6.00 3.00 POINTS 0.00 0.00 2.00 4.00 4.50 5.50 6.00 POINTS 2.00	
Intermittent straight catheter □ Uncontrolled incontinence (frequent linen change), ≥ 3 yrs of age □ Ostomy care - at least daily □ TOTAL	3.50 6.00 3.00 POINTS 0.00 0.00 2.00 4.00 4.50 5.00 5.50 6.00 POINTS 2.00 2.00	
Intermittent straight catheter □ Uncontrolled incontinence (frequent linen change), ≥ 3 yrs of age □ Ostomy care - at least daily □ TOTAL SEIZURES (choose one) No seizure activity □ Mild seizures - at least daily, no intervention □ Mod seizures (req min intervention - at least daily) □ Mod seizures (req min intervention - 2 to 4 times per day) □ Mod seizures (req min intervention - ≥ 5 times per day) □ Severe seizures (req IM/IV/Rectal med administration - at least daily) □ Severe seizures (req IM/IV/Rectal med administration - 2 to 4 times per day) □ Severe seizures (req IM/IV/Rectal med administration - ≥ 5 times per day) □ TOTAL THERAPIES/ORTHOTICS/CASTING (choose any that apply) Fractured or casted limb □ Splinting schedule (off/on at least BID) □ Basic ROM (at least Q shift) □	3.50 6.00 3.00 POINTS 0.00 0.00 2.00 4.00 4.50 5.00 5.50 6.00 POINTS 2.00 2.00 2.00	
Intermittent straight catheter □ Uncontrolled incontinence (frequent linen change), ≥ 3 yrs of age □ Ostomy care - at least daily □ TOTAL	3.50 6.00 3.00 POINTS 0.00 0.00 2.00 4.00 4.50 5.00 5.50 6.00 POINTS 2.00 2.00	

Home Health Manual page 3 of 4

Utah Medicaid Provider Manual	Private Duty Nursing Acuity Grid
Division of Health Care Financing	Updated October 2007

WOUND CARE		POINTS	SCORE
Wound Vac □		2.00	
Tround vao		2.00	
(choose one)			
		2.00	
Stage 1-2, wound care at least daily, dressing change other than trach,		2.00	
PEG, or IV site □			
Stage 3-4, multiple wound sites □		3.00	
	TOTAL		
DEDCONAL CARE		DOINTS	CCORE
PERSONAL CARE		<u>POINTS</u>	<u>SCORE</u>
(choose if applicable)			
Requires personal care/hygiene (≥ 4 yrs of age) □		2.00	
	TOTAL		
BEHAVIOR THAT INTERFERES WITH CARE		POINTS	SCORE
No □		0.00	SCOKE
Yes □		1.00	
	TOTAL		
OTHER ISSUES		POINTS	SCORE
Requires isolation		3.00	<u> </u>
	TOTAL	3.00	
	TOTAL		
* Additional points may be provided based on documentation. NOTE:	A maximum	of ten points v	will be
considered for blood draws.			
TOTAL FOR ALL CATEGORIES ON NURSING ACUITY GRID:			
TOTAL FOR ALL CATEGORIES ON NORSING ACOIT FORID.			
* give points for each treatment			
** give points for each blood draw up to a max of 10 pts			
give points for each blood draw up to a max or to plo			
Care Manager completing:		Date:	
· · · · · · · · · · · · · · · · · · ·			

page 4 of 4 Home Health Manual